

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ForageMax[®]

Arylex[®] active

HERBICIDE

ACTIVE CONSTITUENT: 100 g/L HALAUXIFEN as the methyl ester
50 g/L AMINOPYRALID as the triisopropylamine salt

GROUP I HERBICIDE

A suspension concentrate formulation for post-emergent control of broadleaf weeds in forage brassicas including brassicas undersown with grass pastures, spring or autumn sown canola and spot spraying in grass pastures as specified in the Directions for Use.

Pack Sizes: 1L, 5L

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone: *Australia* 13 11 26. *New Zealand* 0800 764 766.

**EMERGENCY RESPONSE
(ALL HOURS)**
RING FROM ANYWHERE IN
AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

SAFETY DIRECTIONS

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

IN A TRANSPORT
EMERGENCY ONLY
DIAL 000
FOR POLICE OR
FIRE BRIGADE

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **FORAGEMAX[®] ARYLEX[®] ACTIVE HERBICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au



™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

SPRAY DRIFT RESTRAINTS

DO NOT apply ForageMax® with spray droplets smaller than a coarse spray droplet size category according to the “APVMA Compliance Instructions for Mandatory COARSE Droplet Size Categories” located under this title in the **GENERAL INSTRUCTIONS** section of this label.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST** make an accurate written record of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are:

1. Date with start and finish times of application;
2. Location address and paddock/s sprayed;
3. Full name of this product;
4. Amount of product used per hectare and number of hectares applied to;
5. Crop/situation and weed/pest;
6. Wind speed and direction during application;
7. Air temperature and relative humidity during application;
8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;
9. Name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used).

MANDATORY NO SPRAY ZONES

DO NOT apply if there are aquatic or wetland areas including aquacultural ponds, surface streams and rivers within 10 metres downwind from the application area.

DO NOT apply if there are sensitive crops, gardens, landscaping vegetation, protected native vegetation or protected animal habitat within 25 metres downwind from the application area.

DIRECTIONS FOR USE

Canola and Forage brassicas – rape, turnips or swedes including those undersown with grass pasture

RESTRAINTS

DO NOT apply to crops or weeds which may be stressed due to prolonged periods of extreme cold, moisture stress (water-logging or drought) or previous herbicide treatment, as crop effects or reduced levels of weed control may result.

DO NOT sow susceptible crops into paddocks treated the previous season with ForageMax® until after the required plant-back period has elapsed - see **GENERAL INSTRUCTIONS**.

DO NOT spray if rain is likely to occur within three hours or if heavy rain is likely to occur within 48 hours.

DO NOT store mixed spray of ForageMax® in a spray tank for more than 1 day otherwise significant breakdown will occur.

DO NOT use on furrow or flood irrigated crops.

DO NOT apply by air.

Table 1. Canola and Forage brassicas – rape, turnips or swedes including those undersown with grass pastures – boom spray.

READ Crop Safety Directions below

Adjuvant: Always add Uptake® Spraying Oil at 1 L/100 L			
Crop stage: Apply from 4 leaf to 8 leaf brassica crop stage.			
WEED	WEED STAGE	RATE (mL/ha)	CRITICAL COMMENTS
Deadnettle (<i>Lamium amplexicaule</i>)	2 – 6 leaf stage 5-15 cm across	75	Weed growth stage: Treat small weeds up to 6 leaf or 15 cm across for best results. Control will be reduced if larger weeds are treated.
Fumitory (<i>Fumaria</i> spp.)			
Mexican poppy (<i>Argemone mexicana</i>)			
Capeweed (<i>Arctotheca calendula</i>)		100	Conditions: For best results treat weeds that are small, actively growing with excellent soil moisture and not stressed in any way. Control will be reduced if weeds are stressed. Crop stage and type: Only treat brassicas in the 4 - 8 leaf growth stage. Treatment outside these growth stages may result in higher crop injury. DO NOT treat radishes as high injury is likely to result. Canola: For autumn sown canola , apply once at the 4 – 8 leaf stage and avoid treatment in conditions where crop is moderately to severely stressed, due to drought, cold, waterlogging or other factors, as this may result in significant crop effect, which may persist to grain harvest time. Spring-sown canola: apply once at the 4 – 8 leaf growth stage. For spring sown canola that is grazed and left to re-grow a second spray may be applied up the green bud stage (BBCH 51).
Cleavers (<i>Galium aparine</i>)			
Fat hen (<i>Chenopodium album</i>)			
Fleabane (<i>Conyza</i> spp)			
Milk thistle (Sowthistle), (<i>Sonchus oleraceus</i>)			
Rough Sowthistle (<i>Sonchus asper</i>)			
Poppies (<i>Papaver</i> spp.)			
Prickly lettuce (<i>Lactuca serriola</i>)			
Redroot amaranth (<i>Amaranthus retroflexus</i>)			
Shepherd's purse (<i>Capsella bursa-pastoris</i>)			
Small flowered mallow (seedlings) (<i>Malva parviflora</i>)			
Spear thistle (<i>Cirsium arvense</i>)			
Stagger weed (<i>Stachys arvensis</i>)			
Variegated thistle (<i>Silybum marianum</i>)			

Volunteer legumes, including: Chickpea (<i>Cicer arietinum</i>) Clover (<i>Trifolium spp.</i>) Faba bean (<i>Vicia faba</i>) Field pea (<i>Pisum sativum</i>) Lentil (<i>Lens esculentum</i>) Lupin (<i>Lupinus albus</i>) Medics (<i>Medicago spp.</i>) Vetch (<i>Vicia sativa</i>)			Crop effect is less likely when treated in warm conditions, with rapid crop growth and good soil moisture.
---	--	--	---

Table 2. Grass pastures - spot spray (eg knapsack)

Adjuvant: Always add Uptake [®] Spraying Oil at 50 mL/10 L			
WEED	WEED STAGE	RATE (mL/10 L)	CRITICAL COMMENTS
Capeweed (<i>Arctotheca calendula</i>)	2 – 6 leaf stage 5-15cm across	4	Weed growth stage: Treat small weeds up to 6 leaf or 15cm across for best results. Control will be reduced if larger weeds are treated. Conditions: For best results treat weeds that are small, actively growing with excellent soil moisture and not stressed in any way. Control will be reduced if weeds are stressed in any way. Coverage: Spray weeds thoroughly, to completely wet all foliage, to the point of spray runoff.
Milk thistle (Sowthistle), (<i>Sonchus oleraceus</i>)			
Paterson's curse (Salvation Jane) (<i>Echium plantagineum</i>)			
Small flowered mallow (<i>Malva parviflora</i>)			
Spear thistle (<i>Cirsium arvense</i>)			
Stinging nettle (<i>Urtica urens</i>)			
Variegated thistle (<i>Silybum marianum</i>)			

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Grazing/cutting: Forage brassicas, canola or grass pasture: **DO NOT GRAZE OR CUT TREATED CROPS FOR STOCK FEED FOR 2 WEEKS AFTER APPLICATION.**
Harvest (canola): NOT REQUIRED WHEN USED AS DIRECTED

LIVESTOCK DESTINED FOR EXPORT MARKETS

When ForageMax[®] is used as directed and the above withholding period is observed, livestock commodities are considered acceptable for export. However, export requirements are subject to change. Consult your exporter for updated information about specific market requirements.

CROP SAFETY

DO NOT use in forage brassica crops undersown with legume pasture species (eg. medics, clovers), chicory or plantain.

DO NOT use in forage brassica crops undersown with millets if grazing is desirable. ForageMax® will result in significant injury to millets.

DO NOT apply to crops or weeds which may be stressed due to prolonged periods of extreme cold, moisture stress (water-logging or drought) or previous herbicide treatment, as crop effects or reduced levels of weed control may result. When treatment is followed by a severe stress such as drought, prolonged cold, waterlogging or frosty conditions, crop effects such as leaf crinkling, cupping or midvein thickening and whitening may occur. In some cases, there may be an initial growth retardation and biomass reduction. Crops normally recover without loss of dry matter production. Disease, nematode or insect damage after application may also result in crop injury.

GENERAL INSTRUCTIONS

MINIMUM RECROPPING PERIODS following application in forage brassicas

Crop Rotation Recommendations

Use of ForageMax® may prevent early re-establishment of many crops or pastures after treatment. The period that residues persist in the soil will vary according to site conditions such as climate, presence of soil microorganisms, soil temperature and moisture.

Breakdown is fastest in warm, moist soil and slower in cold, dry soil conditions. Land previously treated with ForageMax® should not be rotated to crops other than those listed in the table below. Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas. The ForageMax® treated area may be replanted to any of the specified crops after the interval indicated in the following tables:

Plantback Periods – rain grown

Crops	Rainfall*	Plantback period
Tolerant crop including - Winter cereals (wheat, barley, triticale, oats) or brassica crops (canola, rape or turnips)	> 100 mm	4 months
Ryegrass	> 100mm	2 months
Sensitive crops including (but not limited to) – legumes, linseed, safflower, millet, poppies, potatoes, pyrethrum	>300 mm (with >150 mm Nov – Mar)	12 months

* Rainfall – to safely plant **tolerant crops** (winter cereals or brassicas), there must be sufficient rain and of such distribution to ensure soil wetting to 10 cm for longer than 2 weeks. To safely plant **sensitive crops**, more than 150 mm of the total rain required to safely replant as listed in the table above, must fall over the warmer months of November to March, to ensure soil wetting to depth at least of 10 cm for longer than 2 weeks. Where less than 300 mm rain occurs between application and planting following the sensitive crop, plant a small area of crop before planting whole area, to determine whether safe to plant or not.

Plantback Periods – rain grown with additional irrigation

Crops	Rainfall + irrigation	Plantback period
Sensitive crops including (but not limited to) – legumes, linseed, safflower, millet, poppies, potatoes, pyrethrum	>500 mm (with >250 mm Nov – Mar)	6 months

* Rainfall plus irrigation for 6 month plantback periods – To safely plant **sensitive crops**, more than half of the total rain plus irrigation of 500mm required to safely replant, must fall over the warmer months of November to March, to ensure soil wetting to depth at least of 10 cm for longer than 2-4 weeks. Where less than 500 mm rain plus irrigation occurs between application and planting the following sensitive crop, plant a small area of crop before planting whole area, to determine whether safe to plant or not.

See also PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

Where ForageMax® residue carryover is suspected and sensitive are to be planted, test the treated area as follows:

Where rain allows, plant a small area of the susceptible crop 4 to 6 weeks before desired planting date and take note of any symptoms of injury. If any herbicide symptoms are observed, only plant a winter cereal or brassica crop

Pot bioassay – where not practical to do field bioassay, plant a small number of seeds of the susceptible crop into pots containing soil from the treated field. Do this test 4 to 6 weeks before desired planting date. If any herbicide symptoms are observed, only plant a tolerant crop (see recommendations for northern and southern Australia below).

Clovers are highly sensitive to ForageMax® and therefore are a good test species for a bioassay.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
--------------	----------	------------------

ForageMax® Arylex active Herbicide contains members of the pyridine and arylpicolinate groups of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide. Some naturally occurring weed biotypes resistant to the product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, Corteva Agriscience, consultant or local Department of Agriculture.

MIXING

ForageMax® is a suspension concentrate to be mixed with water. ForageMax® should be added to the spray tank with simultaneous agitation. If ability to agitate the spray tank is limited, pre-mix the ForageMax® in a bucket before adding to the main tank. Once diluted correctly, ForageMax® remains suspended. **THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.**

ForageMax[®] should be mixed and sprayed out within one day. If unable to mix spray immediately, continuous agitation must be maintained to ensure suspension. In tank mixes ForageMax[®] must be in suspension before adding the partner product or Uptake[®] Spraying Oil.

If tank mixing with other products, the following order should be followed:

1. Quarter fill the spray tank, maintaining agitation, then:
2. Add ForageMax[®] (as described above).
3. Add water to half fill the spray tank.
4. Add wettable powders, water dispersible granules or suspension concentrates.
5. Add emulsifiable concentrates.
6. Add Uptake[®] Spraying Oil when spray tank is half full.
7. Add water to bring to the final spray volume.

COMPATIBILITY

ForageMax[®] is compatible with the following:

Insecticides: Success[®] Neo and Lorsban[®] 500 EC.

Adjuvant: Always use Uptake[®] Spraying Oil at 1 L/100 L spray volume for boom spraying or 50 mL/10 L for spot spraying.

APPLICATION

Ground Boom Spraying: Apply ForageMax[®] in 80-200 L water/ha using a coarse spray through accurately calibrated equipment. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping as injury to the crop may occur.

Spot (Dilute) Spraying:

High volume spraying – thorough coverage of foliage to the point of run-off is essential.

Hand-Gun – Apply the recommended mix to give full coverage of leaves and stems through No 6 to No 8 tips at 400 to 500kPa. A spray volume of up to 1000 L per infested hectare of pasture weeds (10 L/100m²) should be used.

Knapsack – Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand-gun. A spray volume of up to 10 L/100m² should be used.

APVMA compliance instructions for mandatory COARSE or larger droplet size categories

Important information

These instructions inform those using this chemical product how to lawfully comply with the requirement of a COARSE or larger spray droplet size category for spray application.

Spray droplet size categories are defined in the ASAE S572 Standard (newer name may also be shown as ASABE) or the BCPC guideline. Nozzle manufacturers may refer to one or both of these documents, to identify droplet size categories; however, for a nozzle to comply with this requirement, the manufacturer must refer to at least one.

Complying with the label requirement to use a specific droplet size category means using the correct nozzle that will deliver that droplet size category under the spray operation conditions being used. The APVMA has approved only the following specific methods for choosing the correct nozzle. Use one of the methods specified

in these instructions to select a correct nozzle to deliver a COARSE or larger droplet size category.

**Instructions for ground application—for COARSE droplet size or larger categories
Mandatory instructions for ground applications**

USE ONLY nozzles that the nozzles' manufacturer has rated to deliver a COARSE, a VERY COARSE or an EXTREMELY COARSE droplet size category, as referenced in ASAE S572 or BCPC. Choose a nozzle that is specified to provide the droplet size category required in the label Spray Drift Restraints.

DO NOT use a higher spray system pressure than the maximum the manufacturer specifies for the selected nozzle to deliver the droplet size category required in the label Spray Drift Restraint.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and their roots and watercourses.

Rinsing: After using ForageMax[®], empty the tank completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain, and clean any filters in the tank, pump, lines, hose and nozzles. After cleaning the tank as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination: Before spraying sensitive crops (see PROTECTION OF CROPS, NATIVES AND OTHER NON-TARGET PLANTS):

Wash the tank and rinse the system as above. Then quarter fill the tank and add a standard alkali based laundry detergent at 500 g (or mL)/100 L water and circulate through the system for at least 15 minutes. If using a concentrated laundry detergent, use 250 g (or mL)/100 L water. Do not use chlorine based cleaners.

Drain the whole system. Remove filter and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Sensitive crops and plants include, but are not limited to chickpeas, clovers, cotton, faba beans, field peas, flowers, fruit trees, hops, lentils, lupins, lucerne, medics, millet, ornamentals, peas, poppies, potatoes, pyrethrum, safflower, shade trees, sub-clover, sugar beet, tobacco, tomatoes, vegetables, vetches, vines (grape and kiwi fruit), wattles and white clover. Field peas, faba beans, lentils and vetches are particularly sensitive

This product will kill or severely injure legumes (clovers, medics) present in the crop at the time of spraying. In the season, following application of this product the regeneration or establishment of sensitive crops may be adversely affected by soil residues.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto non-target vegetation.

DO NOT apply close to or on areas containing roots of desirable vegetation, where treated soil may be washed to areas growing, or to be planted to desirable plants, or on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted to, susceptible crops or plants.

DO NOT move soil, which may have been sprayed, to areas where desirable plants are to be grown.

MANAGEMENT OF RESIDUES IN COMPOST, MULCHES AND ANIMAL WASTE

Do not send treated crops off-farm as fodder or forage. Aminopyralid residues from treated plants may pass into animal manure. If the manure is used to make compost or spread around plants it may cause injury to sensitive plants.

Do not spread manure from animals that have grazed or consumed forage or fodder from treated areas on land used for growing susceptible broadleaf crops.

Dairy and feed pad effluent. Effluent from animals grazing forage brassicas treated with ForageMax[®] within the last four weeks may contain residues. Disposal of this effluent by irrigation during this period may cause damage to clover and other sensitive crops. Effluent from these animals may contain residues for 4 days after removal of the animals from the ForageMax[®] treated crops.

DO NOT send off-farm any effluent (or compost made from it) from animals that have grazed on crops treated with ForageMax[®] within the last 4 weeks until the animals have grazed for 4 days on clean feed. This restriction is not required if 4 weeks has elapsed from treatment to grazing.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

ForageMax[®] is very toxic to aquatic life. **DO NOT** contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the tightly closed original container in a securely locked place, out of direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

DO NOT re-use container.

Containers less than 1 L

Rinse container before disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.

Containers 1 L or greater

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically

marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear appropriate clothing whilst cleaning up small spills (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material and contain in a refuse vessel for disposal. Disposal of the contaminated material must be done in accordance with STATE and/or LOCAL regulations.

APVMA Approval No. 68249/121345

Corteva Agriscience Australia Pty Ltd A.B.N. 24 003 771 659

67 Albert Avenue, Chatswood, NSW 2067

www.corteva.com.au

CUSTOMER SERVICE TOLL FREE

1-800 700 096

TM ® Trademarks of Corteva Agriscience and its affiliated companies. © 2021 Corteva.

Barcode
for stock
identification



This product is GHS compliant. No additional GHS hazard and precautionary statements are required under the Safe Work Australia exemptions for AgVet products.