

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Access[®]

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L TRICLOPYR present as the butoxyethyl ester
120 g/L PICLORAM present as the isooctyl ester

SOLVENT: 389 g/L LIQUID HYDROCARBON

GROUP I HERBICIDE

For selective control of a wide range of woody and noxious weeds in commercial and industrial areas, public lands, fence lines and pastures, by basal bark and cut stump applications as specified in the Directions for Use.

Pack Sizes: 1 L, 2L, 5L 10 L, 20L, 100L

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone: *Australia* 13 11 26. *New Zealand* 0800 764 766.
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

**EMERGENCY RESPONSE
(ALL HOURS)**
RING FROM ANYWHERE IN
AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

SAFETY DIRECTIONS

- Avoid inhaling vapour or spray mist.
- Harmful if swallowed.
- Avoid contact with eyes and skin. Will irritate the eyes, nose, throat and skin.
- Repeated exposure may cause allergic disorders.
- When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrists, a washable hat and elbow-length neoprene gloves and a face shield or goggles.
- If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

IN A TRANSPORT
EMERGENCY ONLY
DIAL 000
FOR POLICE OR
FIRE BRIGADE

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **ACCESS[®] HERBICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au



[™]® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

DIRECTIONS FOR USE**RESTRAINTS:**

- **DO NOT** add water to Access[®] Herbicide.
- **DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.
- **DO NOT** apply to charcoal coated or wet stems as this can repel the diesel mixture.

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /60 L DIESEL²	CRITICAL COMMENTS
Acacia spp. including : Black wattle ¹ Brigalow ¹ Hickory wattle ¹ Mimosa bush Mulga ¹ Sally wattle ¹ Silver wattle ¹	Basal bark method : Plants with stems up to 5 cm basal diameter.	1 L	Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least one metre. Root suckering wattles will need follow-up treatment.
African boxthorn			Treat all stems on multi-stem plants.
Algaroba (<i>Prosopis</i> spp.)	Cut Stump method : Plants up to and in excess of basal bark size. Apply immediately after cut is made.	2 L	This species may become dormant so should not be treated during winter. Regrowth from roots may occur.
<i>Angophora</i> spp. regrowth			
Australian blackthorn			
Bitterbark			
Broadleaf privet			
Brush box			
Cacti including : Common pest pear Harrisia cactus Prickly pear Snake cactus Tiger pear Tree pear			
Calotrope			
Cockspur thorn ¹			
Corkwood wattle			
<i>Corymbia</i> spp. ¹			
Ellangowan poison bush			
<i>Eucalyptus</i> spp. regrowth			
False sandalwood			
Green cestrum			
<i>Grevillea</i> spp.			
Groundsel bush			
Guava		2 L	Regrowth may occur from roots of the main tree.
Hard milkwood		1 L	
Hawthorn			
Heartleaf poison bush			
Honey locust			

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.						
WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /60 L DIESEL²	CRITICAL COMMENTS			
Kiteleaf poison	<p>Basal bark method: Plants with stems up to 5 cm basal diameter.</p> <p>Cut Stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.</p>	1 L	Treat single stem plants only.			
Leucaena ¹						
Limebush						
Lantana						
Narrow leaf hop bush						
Needlewood						
Oleander						
Parkinsonia ¹						
Punty bush						
Red ash (white myrtle)						
Rubbervine						
Swamp box				2 L	Treat single stemmed seedling plants only.	
Sweet briar				1 L	Do not treat during winter.	
Tea trees						
Turpentine						
Turpentine bush						
Whitewood						
Wild orange						Basal bark treatment only.
Wild peach						
Wild Rosemary						
Wild tobacco tree	Cut stump treatment only.					
Wilga						
Yellow bells (<i>Tecoma</i> spp.)	Some root suckering may occur.					
Camphor laurel	<p>Basal bark method: Plants with stems up to 10 cm basal diameter.</p> <p>Cut stump method: Plants in excess of basal bark size. Apply immediately after cut is made.</p>	1 L	Species that become dormant should not be treated during winter. Basal bark method only.			
Peppercorn tree (<i>Schinus molle</i>)						
Prickly acacia						
Sisal hemp				Basal bark method only.		
Tree of Heaven				Species that become dormant should not be treated during winter.		
Willow	4 L	Cut stump method only. Complete control may not occur due to the multi-stem growth of plant and difficulty treating all stems.				
Chinee apple	<p>Basal bark method: Plants with stems up to 15 cm basal diameter.</p> <p>Cut stump method: Plants up to and in excess of basal bark size. Apply immediately after cut is made.</p>	1 L				
<p>¹ The smooth stemmed species <i>Acacia</i> spp. (excluding Mimosa bush), Cockspur thorn, <i>Corymbia</i> spp., Leucaena and Parkinsonia can be treated using the alternative Access[®] ThinLine method for basal bark application. Please refer to the following table.</p> <p>² Biosafe may be used as an alternative carrier to diesel. For alternative biodiesel products, first seek advice from Corteva Agriscience.</p>						

ACCESS® THINLINE BASAL BARK APPLICATION FOR SMOOTH STEMMED SPECIES:

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /10 L DIESEL ²	CRITICAL COMMENTS
<i>Acacia</i> spp. including : Black wattle Brigalow Hickory wattle Mulga Sally wattle Silver wattle <hr/> <i>Cockspur thorn</i> <hr/> <i>Corymbia</i> spp. <hr/> <i>Leucaena</i> <hr/> <i>Parkinsonia</i>	Basal bark method: Plants with stems up to 5 cm basal diameter	1 L	Spray the bark around the stem from ground level up to 5 cm high. Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least one metre. Root suckering wattles will need follow-up treatment.

² Biosafe may be used as an alternative carrier to diesel. For alternative biodiesel products, first seek advice from Corteva Agriscience.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

The use pattern of this product is such that no withholding periods are required.

GENERAL INSTRUCTIONS**MIXING:**

Access® herbicide is designed for use with diesel distillate only. It will NOT mix with water.

Quarter fill the spray unit or mixing container with diesel and add the required amount of Access®. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Only mix sufficient chemical for each day's work, mixing periodically.

Dilution Table:

Rate	Volume of Access® Herbicide			
1L/60L	17 mL	83 mL	167 mL	250 mL
2L/60L	33 mL	167 mL	333 mL	500 mL
4L/60L	67 mL	333 mL	667 mL	1 L
1L/10L	100 mL	500 mL	1 L	1.5 L

Volume of diesel added after Access®	fill to 1 L	fill to 5 L	fill to 10 L	fill to 15 L
---	-------------	-------------	--------------	--------------

COMPATIBILITY:

- Access® is compatible with diesel and Biosafe, and should not be used or diluted with any other chemicals.

For alternative biodiesel products, first seek advice from Corteva Agriscience.

GENERAL APPLICATION:

- Apply only with hand-directed equipment such as a pneumatic sprayer. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers, which use Viton® seals and fittings.
- Spray equipment should be used at low pressures, up to 200 kPa, to avoid splashing.
- Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least 1 metre.
- Species that lose their leaves e.g. sweet briar; or become dormant, e.g. Australian blackthorn or camphor laurel, should not be treated during winter.
- Species with old, rough bark require more thorough wetting than the smooth barked species.
- **DO NOT** apply to charcoal coated or wet stems as this can repel the diesel mixture.

Basal Bark Application Method

- Dilute the indicated volume of Access® Herbicide with an appropriate carrier up to a total volume of 60 L. Liberally spray the bark around the stem from ground level to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).
- Access® ThinLine basal bark application method:
Alternatively, the application band can be reduced to 5 cm high but with the concentration increased proportionally (one part Access® diluted with 9 parts diesel) for the treatment of the smooth stemmed woody weed species: *Acacia* spp. (excluding Mimosa bush), Cockspur thorn, *Corymbia* spp., *Leucaena* and *Parkinsonia*, also indicated with asterisks in the Directions for Use tables.

Cut Stump Method

- Stems should be cut less than 15 cm above ground level.
- **Immediately** apply Access®/carrier mixture **liberally** to the **freshly** cut stump by spraying the cut surface and sides of the stem.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

- After completion of spraying, use a degreaser such as Caltex Kwik-D-Grease® to remove traces of the carrier from the sprayer. Rinse tank and spray through nozzles thoroughly with water to remove degreaser.
After the above, quarter fill the tank with clean water and add a alkali detergent, (e.g. liquid SURF®, OMO®, DRIVE® at 500 mL/100L of water or the powder equivalent at 500 g/100L of water). If using a concentrated laundry detergent use 250 g (or mL)/100 L water. Do not use chlorine based cleaners.
- Shake the water around the sprayer and spray the solution through the nozzles to thoroughly clean the lines and nozzles. Rinse well with clean water to remove detergent.
- To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with water and repeat using detergents (see above).
- **DO NOT** use this equipment for any other purpose.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
-------	---	-----------

Access® Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management the product

is a Group I Herbicide. Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Corteva Agriscience Australia Limited accepts no liability for any losses that may result from the failure of Access® Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Corteva Agriscience representative.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

- DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
- Alongside waterways treat only noxious weeds and poisonous plants.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Treat target weeds only. DO NOT allow mixture to contact non-target species.
- DO NOT use on land to be cultivated for growing sensitive plants. Legumes, vines, vegetables, cotton, tomatoes, ornamentals, coniferous species and many other broadleaf plants are highly susceptible to Access® during both growing and dormant periods.
- Picloram, one of the active ingredients in this product, can remain active in the soil for extended periods depending on soil type, rainfall, temperature, humidity, soil moisture and soil organic matter.
- DO NOT move soil, which has been sprayed, to areas where desirable plants are to be grown.
- DO NOT apply when the treated soil may be washed to areas growing, or to be planted with desirable plants.
- DO NOT apply close to, or on, areas containing roots of desirable plants, or sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted to, susceptible crops or plants.

FOR NATIVE VEGETATION

Use of Access® on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area.

DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

Triple rinse empty containers with diesel before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

5L, 10L, 20L and 100L pack sizes

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

1L pack sizes

Rinse container before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. Dispose of at an approved waste management facility. If no approved waste management facility is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant local, state or territory government regulations. Empty containers and product should not be burnt.

SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.

Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

APVMA Approval No. : 46640/102551

Corteva Agriscience Australia Limited A.B.N. 24 003 771 659

67 Albert Avenue, Chatswood, NSW 2067

www.corteva.com.au

CUSTOMER SERVICE TOLL FREE

1-800 700 096

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © 2021 Corteva.

Barcode
for stock
identification



Hazard and precautionary statements according to classification under GHS (Globally Harmonised System of Classification and Labelling): Combustible liquid. Harmful if swallowed. May be fatal if swallowed and enters airways. Causes skin irritation. May cause an allergic skin reaction. Suspected of causing cancer. May cause damage to organs through prolonged or repeated exposure. Keep away from heat sources. Do not breathe fumes. Wear protective gloves and face protection.